

Sharp Complexity Bounds for Computing Quadrature Formulas for Marginal Distributions of SDEs

Thomas Müller-Gronbach, Uni Passau

joint work with
Klaus Ritter, TU Kaiserslautern
Larisa Yaroslavtseva, Uni Passau

supported by DFG priority program SPP 1324

The Problem

Given a family of d -dimensional systems of SDEs

$$dX_t = a(X_t) dt + b(X_t) dW_t, \quad t \in [0, 1],$$

$$X_0 = x_0 \in \mathbb{R}^d,$$

where $(x_0, a, b) \in \mathcal{H}$.

The Problem

Given a family of d -dimensional systems of SDEs

$$dX_t = a(X_t) dt + b(X_t) dW_t, \quad t \in [0, 1],$$
$$X_0 = x_0 \in \mathbb{R}^d,$$

where $(x_0, a, b) \in \mathcal{H}$.

Approximate

$$S(x_0, a, b) = \mathbb{P}_{X_1}, \quad (x_0, a, b) \in \mathcal{H},$$

by a probability measure with finite support

$$\widehat{S}(x_0, a, b) = \sum_{i=1}^n w_i \cdot \delta_{x_i}.$$

Available: x_0 and finitely many function values of a and b .

The Problem

Given a family of d -dimensional systems of SDEs

$$dX_t = a(X_t) dt + b(X_t) dW_t, \quad t \in [0, 1],$$
$$X_0 = x_0 \in \mathbb{R}^d,$$

where $(x_0, a, b) \in \mathcal{H}$.

Approximate

$$S(x_0, a, b) = \mathbb{P}_{X_1}, \quad (x_0, a, b) \in \mathcal{H},$$

by a probability measure with finite support

$$\widehat{S}(x_0, a, b) = \sum_{i=1}^n w_i \cdot \delta_{x_i}.$$

Available: x_0 and finitely many function values of a and b .

Note that $\widehat{S}(x_0, a, b)$ yields a quadrature formula

$$\int_{\mathbb{R}^d} f d\widehat{S}(x_0, a, b) = \sum_{i=1}^n w_i \cdot f(x_i) \quad \text{for} \quad \int_{\mathbb{R}^d} f dS(x_0, a, b) = \mathbb{E} f(X_1).$$

Complexity of a Computational Task

Given: a class \mathcal{A} of algorithms, error and cost of $A \in \mathcal{A}$.

Complexity of a Computational Task

Given: a class \mathcal{A} of algorithms, error and cost of $A \in \mathcal{A}$.

N -th minimal errors

$$e_N = \inf\{\text{error}(A) : A \in \mathcal{A} \wedge \text{cost}(A) \leq N\}.$$

Complexity of a Computational Task

Given: a class \mathcal{A} of algorithms, error and cost of $A \in \mathcal{A}$.

N -th minimal errors

$$e_N = \inf\{\text{error}(A) : A \in \mathcal{A} \wedge \text{cost}(A) \leq N\}.$$

Enables benchmarks, definition of optimality, analysis of dependence on \mathcal{A} , error, cost.

Complexity of a Computational Task

Given: a class \mathcal{A} of algorithms, error and cost of $A \in \mathcal{A}$.

N -th minimal errors

$$e_N = \inf\{\text{error}(A) : A \in \mathcal{A} \wedge \text{cost}(A) \leq N\}.$$

Enables benchmarks, definition of optimality, analysis of dependence on \mathcal{A} , error, cost.

Typical result $e_N \asymp N^{-\alpha}$

Complexity of a Computational Task

Given: a class \mathcal{A} of algorithms, error and cost of $A \in \mathcal{A}$.

N -th minimal errors

$$e_N = \inf\{\text{error}(A) : A \in \mathcal{A} \wedge \text{cost}(A) \leq N\}.$$

Enables benchmarks, definition of optimality, analysis of dependence on \mathcal{A} , error, cost.

Typical result $e_N \asymp N^{-\alpha}$ consists of

- **upper bound:** $\exists (A_N)_{N \in \mathbb{N}}$ in \mathcal{A} , $c > 0 \forall N \in \mathbb{N}$:

$$\text{cost}(A_N) \leq N \wedge \text{error}(A_N) \leq c \cdot N^{-\alpha}.$$

- **lower bound:** $\exists c > 0 \forall A \in \mathcal{A}$:

$$\text{cost}(A) \leq N \Rightarrow \text{error}(A) \geq c \cdot N^{-\alpha},$$

Complexity of a Computational Task

Given: a class \mathcal{A} of algorithms, error and cost of $A \in \mathcal{A}$.

N -th minimal errors

$$e_N = \inf\{\text{error}(A) : A \in \mathcal{A} \wedge \text{cost}(A) \leq N\}.$$

Enables benchmarks, definition of optimality, analysis of dependence on \mathcal{A} , error, cost.

Typical result $e_N \asymp N^{-\alpha}$ consists of

- **upper bound:** $\exists (A_N)_{N \in \mathbb{N}}$ in \mathcal{A} , $c > 0 \forall N \in \mathbb{N}$:

$$\text{cost}(A_N) \leq N \wedge \text{error}(A_N) \leq c \cdot N^{-\alpha}.$$

- **lower bound:** $\exists c > 0 \forall A \in \mathcal{A}$:

$$\text{cost}(A) \leq N \Rightarrow \text{error}(A) \geq c \cdot N^{-\alpha},$$

Information-based complexity

Traub, Wasilkowski, Woźniakowski (1988), ...

Algorithms, Error and Cost

Deterministic algorithms \hat{S}

- input: $(x_0, a, b) \in \mathcal{H}$,
- use x_0 and function values of a and b via oracle (subroutine),
- exact computation with real numbers and evaluation of elementary functions,
- output:

$$\hat{S}(x_0, a, b) = (n, \underbrace{w_1, \dots, w_n}_{\text{weights } \in [0, 1]}, \underbrace{x_1, \dots, x_n}_{\text{nodes } \in \mathbb{R}^d}) = \sum_{i=1}^n w_i \cdot \delta_{x_i}$$

Algorithms, Error and Cost

Deterministic algorithms \hat{S}

- input: $(x_0, a, b) \in \mathcal{H}$,
- use x_0 and function values of a and b via oracle (subroutine),
- exact computation with real numbers and evaluation of elementary functions,
- output:

$$\hat{S}(x_0, a, b) = (n, \underbrace{w_1, \dots, w_n}_{\text{weights } \in [0, 1]}, \underbrace{x_1, \dots, x_n}_{\text{nodes } \in \mathbb{R}^d}) = \sum_{i=1}^n w_i \cdot \delta_{x_i}$$

Cost of \hat{S}

$$\text{cost}(\hat{S}, (x_0, a, b)) = \# \text{ evaluations of } a, b + \# \text{ instructions}$$

$$\text{cost}(\hat{S}) = \sup_{(x_0, a, b) \in \mathcal{H}} \text{cost}(\hat{S}, (x_0, a, b)).$$

Error of \widehat{S}

$$\text{error}(\widehat{S}, (x_0, a, b)) = \rho_F(S(x_0, a, b), \widehat{S}(x_0, a, b)),$$
$$\text{error}(\widehat{S}) = \sup_{(x_0, a, b) \in \mathcal{H}} \text{error}(\widehat{S}, (x_0, a, b)),$$

where

$$\rho_F(\mu, \widehat{\mu}) = \sup_{f \in F} \left| \int_{\mathbb{R}^d} f \, d\mu - \int_{\mathbb{R}^d} f \, d\widehat{\mu} \right|$$

with a class F of test functions $f : \mathbb{R}^d \rightarrow \mathbb{R}$.

Error of \widehat{S}

$$\text{error}(\widehat{S}, (x_0, a, b)) = \rho_F(S(x_0, a, b), \widehat{S}(x_0, a, b)),$$
$$\text{error}(\widehat{S}) = \sup_{(x_0, a, b) \in \mathcal{H}} \text{error}(\widehat{S}, (x_0, a, b)),$$

where

$$\rho_F(\mu, \widehat{\mu}) = \sup_{f \in F} \left| \int_{\mathbb{R}^d} f \, d\mu - \int_{\mathbb{R}^d} f \, d\widehat{\mu} \right|$$

with a class F of test functions $f : \mathbb{R}^d \rightarrow \mathbb{R}$.

Minimal errors

$$e_N(\mathcal{H}, F) = \inf \{ \text{error}(\widehat{S}) : \text{cost}(\widehat{S}) \leq N \}.$$

Assumptions and Results

Let $s_1, s_2, r \in \mathbb{N}$ and $\beta \geq 0$.

Class of SDEs $\mathcal{H}_{s_1, s_2} = [-1, 1]^d \times H_{s_1}^d \times H_{s_2}^{d \times d}$ with

$$H_s = \{h \in C^s(\mathbb{R}^d; \mathbb{R}) : |h(0)|, \|h^{(\alpha)}\|_\infty \leq 1, 1 \leq |\alpha| \leq s\}.$$

Metric ρ_{F_r} with

$$F_r = \{f \in C^r(\mathbb{R}^d; \mathbb{R}) : |f^{(\alpha)}(x)| \leq 1 + \|x\|^\beta, 1 \leq |\alpha| \leq r\}.$$

Assumptions and Results

Let $s_1, s_2, r \in \mathbb{N}$ and $\beta \geq 0$.

Class of SDEs $\mathcal{H}_{s_1, s_2} = [-1, 1]^d \times H_{s_1}^d \times H_{s_2}^{d \times d}$ with

$$H_s = \{h \in C^s(\mathbb{R}^d; \mathbb{R}) : |h(0)|, \|h^{(\alpha)}\|_\infty \leq 1, 1 \leq |\alpha| \leq s\}.$$

Metric ρ_{F_r} with

$$F_r = \{f \in C^r(\mathbb{R}^d; \mathbb{R}) : |f^{(\alpha)}(x)| \leq 1 + \|x\|^\beta, 1 \leq |\alpha| \leq r\}.$$

Theorem (M-G, Ritter, Yaroslavtseva 2012) For every $\varepsilon > 0$,

$$N^{-\min(s_1, s_2, r)/d} \preceq e_N(\mathcal{H}_{s_1, s_2}, F_r) \preceq N^{-\min(s_1, s_2, r)/d + \varepsilon}.$$

The Lower Bounds

To show:

$$e_N(\mathcal{H}_{s_1, s_2}, \mathcal{F}_r) \geq c \cdot \max(N^{-r/d}, N^{-s_1/d}, N^{-s_2/d}).$$

The Lower Bounds

To show:

$$e_N(\mathcal{H}_{s_1, s_2}, F_r) \geq c \cdot \max(N^{-r/d}, N^{-s_1/d}, N^{-s_2/d}).$$

Use

- $e_N(\mathcal{H}_{s_1, s_2}, F_r) \geq e_N(\mathcal{H}, F)$ if $\mathcal{H} \subset \mathcal{H}_{s_1, s_2}$ and $F \subset F_r$,
- results on minimal errors for weighted integration on \mathbb{R}^d ,
see Wasilkowski, Woźniakowski (2000, 2001).

Case $d = 1$, lower bound N^{-s_1} , see Petras, Ritter (2006): take

$$\mathcal{H} = \{(0, a, 1) : a \in H_{s_1}, \|a\|_\infty \leq N^{-s_1}\}, F = \{f\}$$

with $f \in F_r$, $f \neq \text{const.}$ Then

$$e_N(\mathcal{H}, F) = \inf_{\text{cost}(\widehat{S}) \leq N} \sup_a \left| \int_{\mathbb{R}} f dS(0, a, 1) - \int_{\mathbb{R}} f d\widehat{S}(0, a, 1) \right| .$$

Case $d = 1$, lower bound N^{-s_1} , see Petras, Ritter (2006): take

$$\mathcal{H} = \{(0, a, 1) : a \in H_{s_1}, \|a\|_\infty \leq N^{-s_1}\}, F = \{f\}$$

with $f \in F_r$, $f \neq \text{const.}$ Then

$$e_N(\mathcal{H}, F) = \inf_{\text{cost}(\widehat{S}) \leq N} \sup_a \left| \int_{\mathbb{R}} f dS(0, a, 1) - \int_{\mathbb{R}} f d\widehat{S}(0, a, 1) \right| .$$

We have

$$\int_{\mathbb{R}} f dS(0, a, 1) = \int_{\mathbb{R}} f(x) \cdot \varphi_a(x) dx$$

with a probability density φ_a of $S(0, a, 1)$.

Case $d = 1$, lower bound N^{-s_1} , see Petras, Ritter (2006): take

$$\mathcal{H} = \{(0, a, 1) : a \in H_{s_1}, \|a\|_\infty \leq N^{-s_1}\}, F = \{f\}$$

with $f \in F_r$, $f \neq \text{const.}$ Then

$$e_N(\mathcal{H}, F) = \inf_{\text{cost}(\widehat{S}) \leq N} \sup_a \left| \int_{\mathbb{R}} f dS(0, a, 1) - \int_{\mathbb{R}} f d\widehat{S}(0, a, 1) \right| .$$

We have

$$\int_{\mathbb{R}} f dS(0, a, 1) = \int_{\mathbb{R}} f(x) \cdot \varphi_a(x) dx$$

with a probability density φ_a of $S(0, a, 1)$.

Series expansion of φ_a by means of the parametrix method yields

$$\int_{\mathbb{R}} f(x) \cdot \varphi_a(x) dx = c_f + \underbrace{\int_{\mathbb{R}} a(x) \cdot w_f(x) dx}_{\neq 0} + \underbrace{\sum_{m=2}^{\infty} I_m(f, a^{\otimes m})}_{|\cdot| \leq c_f \cdot N^{-2s_1}} .$$

Case $d = 1$, lower bound N^{-s_1} , see Petras, Ritter (2006): take

$$\mathcal{H} = \{(0, a, 1) : a \in H_{s_1}, \|a\|_\infty \leq N^{-s_1}\}, F = \{f\}$$

with $f \in F_r$, $f \neq \text{const.}$ Then

$$\begin{aligned} e_N(\mathcal{H}, F) &= \inf_{\text{cost}(\widehat{S}) \leq N} \sup_a \left| \int_{\mathbb{R}} f \, dS(0, a, 1) - \int_{\mathbb{R}} f \, d\widehat{S}(0, a, 1) \right| \\ &\geq \inf_{\text{cost}(\widehat{S}) \leq N} \sup_a \left| \int_{\mathbb{R}} a(x) \cdot w_f(x) \, dx + c_f - \int_{\mathbb{R}} f \, d\widehat{S}(0, a, 1) \right| \\ &\quad - c_f \cdot N^{-2s_1} \end{aligned}$$

Case $d = 1$, lower bound N^{-s_1} , see Petras, Ritter (2006): take

$$\mathcal{H} = \{(0, a, 1) : a \in H_{s_1}, \|a\|_\infty \leq N^{-s_1}\}, F = \{f\}$$

with $f \in F_r$, $f \neq \text{const.}$ Then

$$\begin{aligned} e_N(\mathcal{H}, F) &= \inf_{\text{cost}(\widehat{S}) \leq N} \sup_a \left| \int_{\mathbb{R}} f dS(0, a, 1) - \int_{\mathbb{R}} f d\widehat{S}(0, a, 1) \right| \\ &\geq \inf_{\text{cost}(\widehat{S}) \leq N} \sup_a \left| \int_{\mathbb{R}} a(x) \cdot w_f(x) dx + c_f - \int_{\mathbb{R}} f d\widehat{S}(0, a, 1) \right| \\ &\quad - c_f \cdot N^{-2s_1} \\ &\geq \tilde{c}_f \cdot N^{-s_1} - c_f \cdot N^{-2s_1}. \end{aligned}$$

The Upper Bounds

Assume $r = s_1 = s_2 = s$ and $d = 1$.

Construct $\widehat{S}_{n,\varepsilon}$ with

$$\text{cost}(\widehat{S}_{n,\varepsilon}) \leq c \cdot n^{1+\varepsilon} \wedge \text{error}(\widehat{S}_{n,\varepsilon}) \leq c \cdot n^{-s}.$$

The Upper Bounds

Assume $r = s_1 = s_2 = s$ and $d = 1$.

Construct $\widehat{S}_{n,\varepsilon}$ with

$$\text{cost}(\widehat{S}_{n,\varepsilon}) \leq c \cdot n^{1+\varepsilon} \wedge \text{error}(\widehat{S}_{n,\varepsilon}) \leq c \cdot n^{-s}.$$

Idea: use continuity properties of $(x_0, a, b) \mapsto S(x_0, a, b)$.

The Upper Bounds

Assume $r = s_1 = s_2 = s$ and $d = 1$.

Construct $\widehat{S}_{n,\varepsilon}$ with

$$\text{cost}(\widehat{S}_{n,\varepsilon}) \leq c \cdot n^{1+\varepsilon} \wedge \text{error}(\widehat{S}_{n,\varepsilon}) \leq c \cdot n^{-s}.$$

Idea: use continuity properties of $(x_0, a, b) \mapsto S(x_0, a, b)$.

Fix n, ε .

(I) Choose a finite set $\widetilde{\mathcal{H}}$ of SDEs and $\psi: \mathcal{H}_{s,s} \rightarrow \widetilde{\mathcal{H}}$ s.t.

$$\text{cost}(\psi) \leq c \cdot n^{1+\varepsilon} \wedge \rho_{F_1}(S(x_0, a, b), S(\psi(x_0, a, b))) \leq c \cdot n^{-s}.$$

The Upper Bounds

Assume $r = s_1 = s_2 = s$ and $d = 1$.

Construct $\widehat{S}_{n,\varepsilon}$ with

$$\text{cost}(\widehat{S}_{n,\varepsilon}) \leq c \cdot n^{1+\varepsilon} \wedge \text{error}(\widehat{S}_{n,\varepsilon}) \leq c \cdot n^{-s}.$$

Idea: use continuity properties of $(x_0, a, b) \mapsto S(x_0, a, b)$.

Fix n, ε .

(I) Choose a finite set $\tilde{\mathcal{H}}$ of SDEs and $\psi: \mathcal{H}_{s,s} \rightarrow \tilde{\mathcal{H}}$ s.t.

$$\text{cost}(\psi) \leq c \cdot n^{1+\varepsilon} \wedge \rho_{F_1}(S(x_0, a, b), S(\psi(x_0, a, b))) \leq c \cdot n^{-s}.$$


(II) Use approximations $\widehat{S}(\tilde{x}_0, \tilde{a}, \tilde{b})$ for $(\tilde{x}_0, \tilde{a}, \tilde{b}) \in \tilde{\mathcal{H}}$ s.t.

$$|\text{supp}(\widehat{S}(\tilde{x}_0, \tilde{a}, \tilde{b}))| \leq c \cdot n^{1+\varepsilon} \wedge \rho_{F_s}(S(\tilde{x}_0, \tilde{a}, \tilde{b}), \widehat{S}(\tilde{x}_0, \tilde{a}, \tilde{b})) \leq c \cdot n^{-s}.$$

Employ precomputation.


(I) Choose a finite set $\tilde{\mathcal{H}}$ of SDEs and $\psi: \mathcal{H}_{s,s} \rightarrow \tilde{\mathcal{H}}$ s.t.

$$\text{cost}(\psi) \leq c \cdot n^{1+\varepsilon} \wedge \rho_{F_1}(S(x_0, a, b), S(\psi(x_0, a, b))) \leq c \cdot n^{-s}.$$




(I) Choose a finite set $\tilde{\mathcal{H}}$ of SDEs and $\psi: \mathcal{H}_{s,s} \rightarrow \tilde{\mathcal{H}}$ s.t.

$$\text{cost}(\psi) \leq c \cdot n^{1+\varepsilon} \wedge \rho_{F_1}(S(x_0, a, b), S(\psi(x_0, a, b))) \leq c \cdot n^{-s}.$$




(I) Choose a finite set $\tilde{\mathcal{H}}$ of SDEs and $\psi: \mathcal{H}_{s,s} \rightarrow \tilde{\mathcal{H}}$ s.t.

$$\text{cost}(\psi) \leq c \cdot n^{1+\varepsilon} \wedge \rho_{F_1}(S(x_0, a, b), S(\psi(x_0, a, b))) \leq c \cdot n^{-s}.$$




(I) Choose a finite set $\tilde{\mathcal{H}}$ of SDEs and $\psi: \mathcal{H}_{s,s} \rightarrow \tilde{\mathcal{H}}$ s.t.

$$\text{cost}(\psi) \leq c \cdot n^{1+\varepsilon} \wedge \rho_{F_1}(S(x_0, a, b), S(\psi(x_0, a, b))) \leq c \cdot n^{-s}.$$




(I) Choose a finite set $\tilde{\mathcal{H}}$ of SDEs and $\psi: \mathcal{H}_{s,s} \rightarrow \tilde{\mathcal{H}}$ s.t.

$$\text{cost}(\psi) \leq c \cdot n^{1+\varepsilon} \wedge \rho_{F_1}(S(x_0, a, b), S(\psi(x_0, a, b))) \leq c \cdot n^{-s}.$$



(I) Choose a finite set $\tilde{\mathcal{H}}$ of SDEs and $\psi: \mathcal{H}_{s,s} \rightarrow \tilde{\mathcal{H}}$ s.t.

$$\text{cost}(\psi) \leq c \cdot n^{1+\varepsilon} \wedge \rho_{F_1}(S(x_0, a, b), S(\psi(x_0, a, b))) \leq c \cdot n^{-s}.$$




With $n^{1+\varepsilon}$ nodes:

$$\|a - \tilde{a}\|_{[-n^\varepsilon, n^\varepsilon]} \leq c \cdot n^{-s}$$

(I) Choose a finite set $\tilde{\mathcal{H}}$ of SDEs and $\psi: \mathcal{H}_{s,s} \rightarrow \tilde{\mathcal{H}}$ s.t.

$$\text{cost}(\psi) \leq c \cdot n^{1+\varepsilon} \wedge \rho_{F_1}(S(x_0, a, b), S(\psi(x_0, a, b))) \leq c \cdot n^{-s}.$$



With $n^{1+\varepsilon}$ nodes:


$$\|a - \tilde{a}\|_{[-n^{\varepsilon}, n^{\varepsilon}]} \leq c \cdot n^{-s},$$

$$\|b - \tilde{b}\|_{[-n^{\varepsilon}, n^{\varepsilon}]} \leq c \cdot n^{-s},$$

$$\|x_0 - \tilde{x}_0\| \leq c \cdot n^{-s}.$$

(I) Choose a finite set $\tilde{\mathcal{H}}$ of SDEs and $\psi: \mathcal{H}_{s,s} \rightarrow \tilde{\mathcal{H}}$ s.t.

$$\text{cost}(\psi) \leq c \cdot n^{1+\varepsilon} \wedge \rho_{F_1}(S(x_0, a, b), S(\psi(x_0, a, b))) \leq c \cdot n^{-s}.$$



With $n^{1+\varepsilon}$ nodes:

$$\|a - \tilde{a}\|_{[-n^\varepsilon, n^\varepsilon]} \leq c \cdot n^{-s}$$

$$\|b - \tilde{b}\|_{[-n^\varepsilon, n^\varepsilon]} \leq c \cdot n^{-s}$$

$$\|x_0 - \tilde{x}_0\| \leq c \cdot n^{-s}.$$

By a comparison theorem for SDEs:

$$\rho_{F_1}(S(x_0, a, b), S(\tilde{x}_0, \tilde{a}, \tilde{b})) \leq c \cdot n^{-s}.$$

(II) Use approximations $\widehat{S}(\tilde{x}_0, \tilde{a}, \tilde{b})$ for $(\tilde{x}_0, \tilde{a}, \tilde{b}) \in \widetilde{\mathcal{H}}$ s.t.

$$|\text{supp}(\widehat{S}(\tilde{x}_0, \tilde{a}, \tilde{b}))| \leq c \cdot n^{1+\varepsilon} \wedge \rho_{F_s}(S(\tilde{x}_0, \tilde{a}, \tilde{b}), \widehat{S}(\tilde{x}_0, \tilde{a}, \tilde{b})) \leq c \cdot n^{-s}.$$

(II) Use approximations $\widehat{S}(\tilde{x}_0, \tilde{a}, \tilde{b})$ for $(\tilde{x}_0, \tilde{a}, \tilde{b}) \in \widetilde{\mathcal{H}}$ s.t.

$$|\text{supp}(\widehat{S}(\tilde{x}_0, \tilde{a}, \tilde{b}))| \leq c \cdot n^{1+\varepsilon} \wedge \rho_{F_s}(S(\tilde{x}_0, \tilde{a}, \tilde{b}), \widehat{S}(\tilde{x}_0, \tilde{a}, \tilde{b})) \leq c \cdot n^{-s}.$$

a) Iterative quantization of the Euler scheme with step-size $\Delta = n^{-2s}$:

$$\begin{aligned}\widehat{X}_{k+1} &= \widehat{X}_k + \tilde{a}(\widehat{X}_k) \cdot \Delta + \tilde{b}(\widehat{X}_k) \cdot \Delta_k W, \quad k = 0, \dots, n^{2s}, \\ T: \mathbb{R} &\rightarrow \mathbb{R}, \text{ with appropriate finite range } T(\mathbb{R}),\end{aligned}$$

$$\begin{aligned}\overline{X}_{k+1} &= \tilde{X}_k + \tilde{a}(\tilde{X}_k) \cdot \Delta + \tilde{b}(\tilde{X}_k) \cdot \Delta_k W, \\ \tilde{X}_{k+1} &= T(\overline{X}_{k+1}).\end{aligned}$$

Yields a measure $Q = P_{\tilde{X}_{n^{2s}}}$ with

$$|\text{supp}(Q)| \leq c \cdot n^{3s} \quad \wedge \quad \rho_{F_1}(S(\tilde{x}_0, \tilde{a}, \tilde{b}), Q) \leq c \cdot n^{-s}.$$

(II) Use approximations $\widehat{S}(\tilde{x}_0, \tilde{a}, \tilde{b})$ for $(\tilde{x}_0, \tilde{a}, \tilde{b}) \in \widetilde{\mathcal{H}}$ s.t.

$$|\text{supp}(\widehat{S}(\tilde{x}_0, \tilde{a}, \tilde{b}))| \leq c \cdot n^{1+\varepsilon} \wedge \rho_{F_s}(S(\tilde{x}_0, \tilde{a}, \tilde{b}), \widehat{S}(\tilde{x}_0, \tilde{a}, \tilde{b})) \leq c \cdot n^{-s}.$$

a) Iterative quantization of the Euler scheme with step-size $\Delta = n^{-2s}$:

$$\begin{aligned}\widehat{X}_{k+1} &= \widehat{X}_k + \tilde{a}(\widehat{X}_k) \cdot \Delta + \tilde{b}(\widehat{X}_k) \cdot \Delta_k W, \quad k = 0, \dots, n^{2s}, \\ T: \mathbb{R} &\rightarrow \mathbb{R}, \text{ with appropriate finite range } T(\mathbb{R}),\end{aligned}$$

$$\begin{aligned}\overline{X}_{k+1} &= \tilde{X}_k + \tilde{a}(\tilde{X}_k) \cdot \Delta + \tilde{b}(\tilde{X}_k) \cdot \Delta_k W, \\ \tilde{X}_{k+1} &= T(\overline{X}_{k+1}).\end{aligned}$$

Yields a measure $Q = P_{\tilde{X}_{n^{2s}}}$ with

$$|\text{supp}(Q)| \leq c \cdot n^{3s} \quad \wedge \quad \rho_{F_1}(S(\tilde{x}_0, \tilde{a}, \tilde{b}), Q) \leq c \cdot n^{-s}.$$

b) Apply a support reduction algorithm to Q , see Davis (1967).

Yields a measure $\widehat{S}(\tilde{x}_0, \tilde{a}, \tilde{b})$ with

$$|\text{supp}(\widehat{S}(\tilde{x}_0, \tilde{a}, \tilde{b}))| \leq c \cdot n^{1+\varepsilon} \quad \wedge \quad \rho_{F_s}(Q, \widehat{S}(\tilde{x}_0, \tilde{a}, \tilde{b})) \leq c \cdot n^{-s}.$$

Discussion

- We have

$$e_N(\mathcal{H}_{s_1, s_2}, F_r) \asymp N^{-\min(r, s_1, s_2)/d + \varepsilon}.$$

Discussion

- We have

$$e_N(\mathcal{H}_{s_1, s_2}, F_r) \asymp N^{-\min(r, s_1, s_2)/d + \varepsilon}.$$

- Upper bound by algorithms $\widehat{S}_{\varepsilon, n}$ with

$$\text{cost}(\widehat{S}_{\varepsilon, n}) \preceq n^{d(1+\varepsilon)} \quad \wedge \quad \text{error}(\widehat{S}_{\varepsilon, n}) \preceq n^{-\min(r, s_1, s_2)}.$$

Discussion

- We have

$$e_N(\mathcal{H}_{s_1, s_2}, F_r) \asymp N^{-\min(r, s_1, s_2)/d + \varepsilon}.$$

- Upper bound by algorithms $\widehat{S}_{\varepsilon, n}$ with

$$\text{cost}(\widehat{S}_{\varepsilon, n}) \preceq n^{d(1+\varepsilon)} \quad \wedge \quad \text{error}(\widehat{S}_{\varepsilon, n}) \preceq n^{-\min(r, s_1, s_2)}.$$

Extensive precomputations: roughly, $n^{d \cdot \min(s_1, s_2) \cdot n}$ operations.

Discussion

- We have

$$e_N(\mathcal{H}_{s_1, s_2}, F_r) \asymp N^{-\min(r, s_1, s_2)/d + \varepsilon}.$$

- Upper bound by algorithms $\widehat{S}_{\varepsilon, n}$ with

$$\text{cost}(\widehat{S}_{\varepsilon, n}) \preceq n^{d(1+\varepsilon)} \quad \wedge \quad \text{error}(\widehat{S}_{\varepsilon, n}) \preceq n^{-\min(r, s_1, s_2)}.$$

Extensive precomputations: roughly, $n^{d \cdot \min(s_1, s_2) \cdot n}$ operations.

- Implementable modification $\widetilde{S}_{\varepsilon, n}$ of $\widehat{S}_{\varepsilon, n}$ with

$$\text{error}(\widetilde{S}_{\varepsilon, n}) \preceq (\text{cost}(\widetilde{S}_{\varepsilon, n}))^{-r/(3dr+2r+3d)+\varepsilon}.$$

- Cubature on Wiener space

Kusuoka (2001), . . . , Lyons, Victoir (2004), Litterer, Lyons (2010).

Algorithm $\widehat{S}_{m,n}$ based on

- a) Quadrature formula on the Wiener space with nodes given by bv-paths; exact for multiple Stratonovich integrals up to order m .
- b) Nonuniform time discretization $0 = t_0 < \dots < t_n = 1$

Given the discrete approximation μ_ℓ of $P_{X_{t_\ell}}$, solve the SDE along the scaled bv-paths on subinterval $[t_\ell, t_{\ell+1}]$ and recombine to get $\mu_{\ell+1}$.

- Cubature on Wiener space

Kusuoka (2001), . . . , Lyons, Victoir (2004), Litterer, Lyons (2010).

Algorithm $\widehat{S}_{m,n}$ based on

- a) Quadrature formula on the Wiener space with nodes given by bv-paths; exact for multiple Stratonovich integrals up to order m .
- b) Nonuniform time discretization $0 = t_0 < \dots < t_n = 1$

Given the discrete approximation μ_ℓ of $P_{X_{t_\ell}}$, solve the SDE along the scaled bv-paths on subinterval $[t_\ell, t_{\ell+1}]$ and recombine to get $\mu_{\ell+1}$.

Class of SDEs: $x_0 \in \mathbb{R}^d$, $a_i, b_{i,j} \in C_b^\infty$, uniform Hörmander condition.

Metric: ρ_F with $F = \text{Lip}_1(\mathbb{R}^d; \mathbb{R})$.

- Cubature on Wiener space

Kusuoka (2001), . . . , Lyons, Victoir (2004), Litterer, Lyons (2010).

Algorithm $\widehat{S}_{m,n}$ based on

- a) Quadrature formula on the Wiener space with nodes given by bv-paths; exact for multiple Stratonovich integrals up to order m .
- b) Nonuniform time discretization $0 = t_0 < \dots < t_n = 1$

Given the discrete approximation μ_ℓ of $P_{X_{t_\ell}}$, solve the SDE along the scaled bv-paths on subinterval $[t_\ell, t_{\ell+1}]$ and recombine to get $\mu_{\ell+1}$.

Class of SDEs: $x_0 \in \mathbb{R}^d$, $a_i, b_{i,j} \in C_b^\infty$, uniform Hörmander condition.

Metric: ρ_F with $F = \text{Lip}_1(\mathbb{R}^d; \mathbb{R})$.

Then

$$\text{error}(\widehat{S}_{m,n}) \preceq \text{cost}(\widehat{S}_{m,n})^{-\frac{m-1}{dm} + \varepsilon}.$$

- Cubature on Wiener space

Kusuoka (2001), . . . , Lyons, Victoir (2004), Litterer, Lyons (2010).

Algorithm $\widehat{S}_{m,n}$ based on

- Quadrature formula on the Wiener space with nodes given by bv-paths; exact for multiple Stratonovich integrals up to order m .
- Nonuniform time discretization $0 = t_0 < \dots < t_n = 1$

Given the discrete approximation μ_ℓ of $P_{X_{t_\ell}}$, solve the SDE along the scaled bv-paths on subinterval $[t_\ell, t_{\ell+1}]$ and recombine to get $\mu_{\ell+1}$.

Class of SDEs: $x_0 \in \mathbb{R}^d$, $a_i, b_{i,j} \in C_b^\infty$, uniform Hörmander condition.

Metric: ρ_F with $F = \text{Lip}_1(\mathbb{R}^d; \mathbb{R})$.

Then

$$\text{error}(\widehat{S}_{m,n}) \preceq \text{cost}(\widehat{S}_{m,n})^{-\frac{m-1}{dm} + \varepsilon}.$$

For $a_i, b_{i,j} \in C_b^s$, $s \geq 2$:

$$\text{error}(\widehat{S}_{s-1,n}) \preceq \text{cost}(\widehat{S}_{s-1,n})^{-\frac{s-2}{d(s-1)} + \varepsilon}.$$

- Cubature on Wiener space

Kusuoka (2001), . . . , Lyons, Victoir (2004), Litterer, Lyons (2010).

Class of SDEs: $x_0 \in \mathbb{R}^d$, $a_i, b_{i,j} \in C_b^s$, $s \geq 2$, unif. Hörmander condition.

Metric: ρ_F with $F = \text{Lip}_1(\mathbb{R}^d; \mathbb{R})$.

Algorithms \widehat{S}_n with

$$\text{error}(\widehat{S}_n) \preceq \text{cost}(\widehat{S}_n)^{-\frac{s-2}{d(s-1)} + \varepsilon}.$$

- Cubature on Wiener space

Kusuoka (2001), . . . , Lyons, Victoir (2004), Litterer, Lyons (2010).

Class of SDEs: $x_0 \in \mathbb{R}^d$, $a_i, b_{i,j} \in C_b^s$, $s \geq 2$, unif. Hörmander condition.

Metric: ρ_F with $F = \text{Lip}_1(\mathbb{R}^d; \mathbb{R})$.

Algorithms \widehat{S}_n with

$$\text{error}(\widehat{S}_n) \preceq \text{cost}(\widehat{S}_n)^{-\frac{s-2}{d(s-1)} + \varepsilon}.$$

- Sparse Markov chain based on

Wagner-Platen scheme with non-uniform time steps

M-G, Yaroslavtseva (2012)

Class of SDEs: $d = 1$, $x_0 \in [-1, 1]$, $a, b \in C_b^s$ with $s \geq 6$, $|b| \geq \delta$.

Metric: ρ_F with $F = \text{Lip}_1(\mathbb{R}; \mathbb{R})$.

Algorithms \widehat{S}_n with

$$\text{error}(\widehat{S}_n) \preceq \text{cost}(\widehat{S}_n)^{-(1-\varepsilon)}.$$